

United States Department of Agriculture

Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Union County, Pennsylvania

Gordon Farm



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

| Preface Soil Map | |
|---|----|
| Soil Map | 6 |
| Legend | 7 |
| Map Unit Legend | |
| Map Unit Descriptions | 8 |
| Union County, Pennsylvania | 10 |
| AbB—Albrights silt loam, 3 to 8 percent slopes | 10 |
| CaB—Calvin-Klinesville shaly silt loams, 3 to 8 percent slopes | 11 |
| CaC—Calvin-Klinesville shaly silt loams, 8 to 15 percent slopes | 13 |
| Hv—Holly silt loam | 15 |
| MkB—Meckesville silt loam, 3 to 8 percent slopes | 16 |
| MkC—Meckesville silt loam, 8 to 15 percent slopes | 17 |
| ShB—Shelmadine silt loam, 3 to 8 percent slopes | 18 |
| W—Water | 20 |
| WkE—Weikert and Klinesville shaly silt loams, steep | 20 |

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



| | MAP LEGEND | | | MAP INFORMATION | |
|-------------|---|------------------|----------------------------------|---|--|
| | terest (AOI) Area of Interest (AOI) | 8 | Spoil Area Stony Spot | The soil surveys that comprise your AOI were mapped at 1:20,000. | |
| Soils | Soil Map Unit Polygons | 00 V | Very Stony Spot Wet Spot | Warning: Soil Map may not be valid at this scale. | |
| Ĩ | Soil Map Unit Lines Soil Map Unit Points | ۵ | Other Special Line Features | Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of | |
| ဖ | ÷ | | tures Streams and Canals | contrasting soils that could have been shown at a more detailed scale. | |
| × | Clay Spot Closed Depression | Transport +++ | ation Rails | Please rely on the bar scale on each map sheet for map measurements. | |
| ◇ ¥ | Gravel Pit Gravelly Spot | ~ | Interstate Highways US Routes | Source of Map: Natural Resources Conservation Service Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857) | |
| Ø | Landfill Lava Flow | ~ | Major Roads Local Roads | Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts | |
| 人 小 の | Marsh or swamp Mine or Quarry | Backgrou | Aerial Photography | distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required. | |
| 0 | Miscellaneous Water Perennial Water | | | This product is generated from the USDA-NRCS certified data as of the version date(s) listed below. | |
| ~ + | Rock Outcrop Saline Spot | | | Soil Survey Area: Union County, Pennsylvania Survey Area Data: Version 17, Sep 7, 2023 | |
| .: | Sandy Spot Severely Eroded Spot | | | Soil map units are labeled (as space allows) for map scales 1:50,000 or larger. | |
| ♦ ≥ | Sinkhole Slide or Slip | | | Date(s) aerial images were photographed: Jul 6, 2020—Nov 7, 2020 | |
| ø | Sodic Spot | | | The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident. | |

| Map Unit Symbol | Map Unit Name | Acres in AOI | Percent of AOI | | |
|-----------------------------|--|---------------------------------------|----------------|--|--|
| AbB | Albrights silt loam, 3 to 8 percent slopes | 6.8 | 14.7% | | |
| СаВ | Calvin-Klinesville shaly silt loams, 3 to 8 percent slopes | 3.9 | 8.4% | | |
| CaC | Calvin-Klinesville shaly silt loams, 8 to 15 percent slopes | , , , , , , , , , , , , , , , , , , , | | | |
| Hv | Holly silt loam | am 9.4 | | | |
| MkB | Meckesville silt loam, 3 to 8 percent slopes | 2.7 | 5.9% | | |
| MkC | Meckesville silt loam, 8 to 15 percent slopes | 8.6 | 18.6% | | |
| ShB | Shelmadine silt loam, 3 to 8 percent slopes | 2.6 | 5.7% | | |
| W | Water | 0.8 | 1.7% | | |
| WkE | Weikert and Klinesville shaly silt loams, steep | 2.0 | 4.3% | | |
| Totals for Area of Interest | | 46.3 | 100.0% | | |

Map Unit Legend

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas

are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Union County, Pennsylvania

AbB—Albrights silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 14tj Elevation: 500 to 2,800 feet Mean annual precipitation: 34 to 48 inches Mean annual air temperature: 40 to 60 degrees F Frost-free period: 130 to 220 days Farmland classification: All areas are prime farmland

Map Unit Composition

Albrights and similar soils: 80 percent *Minor components:* 20 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Albrights

Setting

Landform: Ridges Landform position (two-dimensional): Toeslope, footslope Landform position (three-dimensional): Head slope Down-slope shape: Convex Across-slope shape: Concave Parent material: Fine-loamy colluvium derived from sedimentary rock

Typical profile

H1 - 0 to 10 inches: silt loam *H2 - 10 to 30 inches:* clay loam *H3 - 30 to 60 inches:* silt loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 18 to 32 inches to fragipan
Drainage class: Moderately well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)
Depth to water table: About 12 to 28 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 2.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 2e Hydrologic Soil Group: C/D Ecological site: F147XY002PA - Mixed Sedimentary Upland Hydric soil rating: No

Minor Components

Leck kill

Percent of map unit: 5 percent Hydric soil rating: No

Shelmadine

Percent of map unit: 5 percent Landform: Drainageways Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope Down-slope shape: Concave Across-slope shape: Concave Hydric soil rating: Yes

Meckesville

Percent of map unit: 5 percent Landform: Mountain valleys Landform position (two-dimensional): Footslope Landform position (three-dimensional): Lower third of mountainflank Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

Alvira

Percent of map unit: 5 percent Landform: Hillslopes Landform position (two-dimensional): Footslope Landform position (three-dimensional): Interfluve Down-slope shape: Concave Across-slope shape: Concave Hydric soil rating: No

CaB—Calvin-Klinesville shaly silt loams, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 14v5 Elevation: 300 to 1,600 feet Mean annual precipitation: 36 to 50 inches Mean annual air temperature: 45 to 57 degrees F Frost-free period: 120 to 217 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Calvin and similar soils: 50 percent *Klinesville and similar soils:* 30 percent *Minor components:* 20 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Calvin

Setting

Landform: Hillslopes Landform position (three-dimensional): Side slope, crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from siltstone

Typical profile

H1 - 0 to 8 inches: channery silt loam H2 - 8 to 25 inches: very channery silt loam H3 - 25 to 30 inches: very channery silt loam

R - 30 to 34 inches: unweathered bedrock

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 20 to 40 inches to lithic bedrock
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 2e Hydrologic Soil Group: B Ecological site: F147XY008PA - Shallow Mixed Sedimentary Upland Hydric soil rating: No

Description of Klinesville

Setting

Landform: Valleys, ridges Landform position (two-dimensional): Backslope, shoulder Landform position (three-dimensional): Side slope Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from siltstone

Typical profile

H1 - 0 to 7 inches: channery silt loam
H2 - 7 to 11 inches: very channery silt loam
H3 - 11 to 15 inches: very channery silt loam
R - 15 to 19 inches: unweathered bedrock

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Drainage class: Somewhat excessively drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: D *Ecological site:* F147XY008PA - Shallow Mixed Sedimentary Upland *Hydric soil rating:* No

Minor Components

Leck kill

Percent of map unit: 10 percent *Hydric soil rating:* No

Weikert

Percent of map unit: 5 percent Hydric soil rating: No

Berks

Percent of map unit: 5 percent Hydric soil rating: No

CaC—Calvin-Klinesville shaly silt loams, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: I4v6 Elevation: 300 to 1,600 feet Mean annual precipitation: 36 to 50 inches Mean annual air temperature: 45 to 57 degrees F Frost-free period: 120 to 217 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Calvin and similar soils: 50 percent Klinesville and similar soils: 30 percent Minor components: 20 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Calvin

Setting

Landform: Hillslopes Landform position (three-dimensional): Side slope, crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from siltstone

Typical profile

H1 - 0 to 8 inches: channery silt loam
H2 - 8 to 25 inches: very channery silt loam
H3 - 25 to 30 inches: very channery silt loam
R - 30 to 34 inches: unweathered bedrock

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock Drainage class: Well drained Runoff class: Medium Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr) Depth to water table: More than 80 inches Frequency of flooding: None Frequency of ponding: None Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: F147XY008PA - Shallow Mixed Sedimentary Upland Hydric soil rating: No

Description of Klinesville

Setting

Landform: Valleys, ridges Landform position (two-dimensional): Backslope, shoulder Landform position (three-dimensional): Side slope Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from siltstone

Typical profile

H1 - 0 to 7 inches: channery silt loam
H2 - 7 to 11 inches: very channery silt loam
H3 - 11 to 15 inches: very channery silt loam
R - 15 to 19 inches: unweathered bedrock

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Drainage class: Somewhat excessively drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 4e Hydrologic Soil Group: D Ecological site: F147XY008PA - Shallow Mixed Sedimentary Upland Hydric soil rating: No

Minor Components

Leck kill

Percent of map unit: 10 percent Hydric soil rating: No

Berks

Percent of map unit: 5 percent *Hydric soil rating:* No

Weikert

Percent of map unit: 5 percent *Hydric soil rating:* No

Hv—Holly silt loam

Map Unit Setting

National map unit symbol: I4w1 Elevation: 400 to 1,170 feet Mean annual precipitation: 30 to 45 inches Mean annual air temperature: 45 to 54 degrees F Frost-free period: 120 to 187 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Holly and similar soils: 90 percent Minor components: 10 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Holly

Setting

Landform: Depressions on flood plains, backswamps Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope Down-slope shape: Concave Across-slope shape: Linear Parent material: Loamy alluvium derived from sandstone and shale

Typical profile

H1 - 0 to 11 inches: silt loam
H2 - 11 to 42 inches: silty clay loam
H3 - 42 to 60 inches: stratified very gravelly loamy sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 2.00 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Available water supply, 0 to 60 inches: High (about 11.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 4w Hydrologic Soil Group: B/D Ecological site: F147XY011PA - Poorly Drained Fine Mixed Floodplain Hydric soil rating: Yes

Minor Components

Basher

Percent of map unit: 5 percent Hydric soil rating: No

Holly, ponded

Percent of map unit: 5 percent Landform: Flood plains Hydric soil rating: Yes

MkB—Meckesville silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 14wg Elevation: 400 to 2,800 feet Mean annual precipitation: 34 to 48 inches Mean annual air temperature: 45 to 55 degrees F Frost-free period: 120 to 220 days Farmland classification: All areas are prime farmland

Map Unit Composition

Meckesville and similar soils: 85 percent *Minor components:* 15 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Meckesville

Setting

Landform: Mountain valleys, mountain slopes Landform position (two-dimensional): Footslope Landform position (three-dimensional): Lower third of mountainflank Down-slope shape: Concave Across-slope shape: Linear Parent material: Sandstone, siltstone and shale colluvium derived from sedimentary rock

Typical profile

- H1 0 to 4 inches: silt loam
- H2 4 to 36 inches: silt loam
- H3 36 to 60 inches: gravelly silty clay loam

Properties and qualities

Slope: 3 to 8 percent

Custom Soil Resource Report

Depth to restrictive feature: 25 to 48 inches to fragipan Drainage class: Well drained Runoff class: Medium Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr) Depth to water table: About 28 to 48 inches Frequency of flooding: None Frequency of ponding: None Available water supply, 0 to 60 inches: Low (about 4.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 2e Hydrologic Soil Group: C Ecological site: F147XY002PA - Mixed Sedimentary Upland Hydric soil rating: No

Minor Components

Albrights

Percent of map unit: 5 percent Hydric soil rating: No

Leck kill

Percent of map unit: 5 percent Hydric soil rating: No

Calvin

Percent of map unit: 5 percent *Hydric soil rating:* No

MkC—Meckesville silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: I4wh Elevation: 400 to 2,800 feet Mean annual precipitation: 34 to 48 inches Mean annual air temperature: 45 to 55 degrees F Frost-free period: 120 to 190 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Meckesville and similar soils: 85 percent *Minor components:* 15 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Meckesville

Setting

Landform: Mountain valleys, mountain slopes Landform position (two-dimensional): Footslope Landform position (three-dimensional): Lower third of mountainflank Down-slope shape: Concave Across-slope shape: Linear Parent material: Sandstone, siltstone and shale colluvium derived from sedimentary rock

Typical profile

H1 - 0 to 4 inches: silt loam
H2 - 4 to 36 inches: silt loam
H3 - 36 to 60 inches: gravelly silty clay loam

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 25 to 48 inches to fragipan
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)
Depth to water table: About 28 to 48 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 4.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: C Ecological site: F147XY002PA - Mixed Sedimentary Upland Hydric soil rating: No

Minor Components

Calvin

Percent of map unit: 5 percent Hydric soil rating: No

Allenwood

Percent of map unit: 5 percent Hydric soil rating: No

Leck kill

Percent of map unit: 5 percent Hydric soil rating: No

ShB—Shelmadine silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: I4wv Elevation: 430 to 1,850 feet Mean annual precipitation: 36 to 46 inches Mean annual air temperature: 40 to 60 degrees F Frost-free period: 130 to 220 days Farmland classification: Not prime farmland

Map Unit Composition

Shelmadine and similar soils: 80 percent Minor components: 20 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Shelmadine

Setting

Landform: Drainageways Down-slope shape: Concave Across-slope shape: Concave Parent material: Loamy till

Typical profile

H1 - 0 to 7 inches: silt loam H2 - 7 to 24 inches: silty clay loam H3 - 24 to 50 inches: channery loam H4 - 50 to 70 inches: channery loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 18 to 30 inches to fragipan
Drainage class: Poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 0 to 6 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 3.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 4w Hydrologic Soil Group: C/D Ecological site: F147XY011PA - Poorly Drained Fine Mixed Floodplain Hydric soil rating: Yes

Minor Components

Watson

Percent of map unit: 10 percent Hydric soil rating: No

Alvira

Percent of map unit: 5 percent Landform: Hillslopes Landform position (two-dimensional): Footslope Landform position (three-dimensional): Interfluve Down-slope shape: Concave Across-slope shape: Concave Hydric soil rating: No

Albrights

Percent of map unit: 5 percent Hydric soil rating: No

W-Water

Map Unit Setting

National map unit symbol: ktpj Mean annual precipitation: 36 to 46 inches Mean annual air temperature: 44 to 57 degrees F Frost-free period: 130 to 180 days Farmland classification: Not prime farmland

Map Unit Composition

Water: 100 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

WkE—Weikert and Klinesville shaly silt loams, steep

Map Unit Setting

National map unit symbol: I4x8 Elevation: 300 to 1,600 feet Mean annual precipitation: 36 to 50 inches Mean annual air temperature: 45 to 57 degrees F Frost-free period: 120 to 217 days Farmland classification: Not prime farmland

Map Unit Composition

Weikert and similar soils: 40 percent Klinesville and similar soils: 30 percent Minor components: 30 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Weikert

Setting

Landform: Hills Landform position (two-dimensional): Backslope, shoulder Landform position (three-dimensional): Side slope, crest Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from shale and siltstone

Typical profile

H1 - 0 to 7 inches: channery silt loam *H2 - 7 to 15 inches:* very channery silt loam *H3 - 15 to 19 inches:* bedrock

Properties and qualities

Slope: 25 to 75 percent *Depth to restrictive feature:* 10 to 20 inches to lithic bedrock Drainage class: Somewhat excessively drained Runoff class: High Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr) Depth to water table: More than 80 inches Frequency of flooding: None Frequency of ponding: None Available water supply, 0 to 60 inches: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 7e Hydrologic Soil Group: D Ecological site: F147XY008PA - Shallow Mixed Sedimentary Upland Hydric soil rating: No

Description of Klinesville

Setting

Landform: Valleys, ridges Landform position (two-dimensional): Backslope, shoulder Landform position (three-dimensional): Side slope Down-slope shape: Convex Across-slope shape: Convex Parent material: Residuum weathered from siltstone

Typical profile

H1 - 0 to 7 inches: channery silt loam
H2 - 7 to 11 inches: very channery silt loam
H3 - 11 to 15 inches: very channery silt loam
R - 15 to 19 inches: bedrock

Properties and qualities

Slope: 25 to 75 percent
Depth to restrictive feature: 10 to 20 inches to lithic bedrock
Drainage class: Somewhat excessively drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 7e Hydrologic Soil Group: D Ecological site: F147XY008PA - Shallow Mixed Sedimentary Upland Hydric soil rating: No

Minor Components

Hartleton

Percent of map unit: 10 percent Landform: — error in exists on — Landform position (two-dimensional): Shoulder, backslope Landform position (three-dimensional): Side slope Down-slope shape: Concave, linear Across-slope shape: Linear, concave Hydric soil rating: No

Berks

Percent of map unit: 10 percent Hydric soil rating: No

Leck kill

Percent of map unit: 5 percent Hydric soil rating: No

Rushtown

Percent of map unit: 5 percent *Hydric soil rating:* No